

Confidential



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

SENIOR CERTIFICATE EXAMINATIONS/ NATIONAL SENIOR CERTIFICATE EXAMINATIONS

TECHNICAL SCIENCES P2

MAY/JUNE 2024

MARKS: 75

TIME: 1½ hours

This question paper consists of 9 pages and 4 data sheets.



INSTRUCTIONS AND INFORMATION

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of SIX questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You are advised to use the attached DATA SHEETS.
8. Round off your FINAL numerical answers to a minimum of TWO decimal places.
9. Give brief motivations, discussions, etc. where required.
10. Write neatly and legibly.



QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.5) in the ANSWER BOOK, e.g. 1.6 D.

- 1.1 The name of the functional group of ETHANOL is ...
A carboxyl.
B carbonyl.
C hydroxyl.
D formyl. (2)
- 1.2 Which homologous series has London forces, dipole-dipole forces and hydrogen bonds?
A Alkanes
B Ketones
C Aldehydes
D Carboxylic acids (2)
- 1.3 An element in group 3 that can be used for doping:
A Germanium
B Gallium
C Silicon
D Tin (2)
- 1.4 Which ONE of the following combinations of standard conditions is applicable to a galvanic cell with non-gaseous reactants and products?
A $1 \text{ mol}\cdot\text{dm}^{-3}$; 101,3 kPa; 25 K
B $1 \text{ mol}\cdot\text{dm}^{-3}$; 101,3 kPa
C $1 \text{ mol}\cdot\text{dm}^{-3}$; 298 K
D $1 \text{ mol}\cdot\text{dm}^{-3}$; 0 K (2)
- 1.5 A solution that can conduct an electric current through the movement of ions:
A Oxidising agent
B Reducing agent
C Electrolysis
D Electrolyte (2)

[10]

QUESTION 2 (Start on a new page.)

The table below represents six organic molecules with different functional groups.

A	$\text{H}-\text{C}\equiv\text{C}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\text{H}$	B	$\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{C}-\text{O}-\text{H} \end{array}$
C	$\begin{array}{c} \text{H} & & \text{O} & \text{H} \\ & & & \\ \text{H}-\text{C}-\text{O}-\text{C}-\text{C}-\text{H} \\ & & & \\ \text{H} & & & \text{H} \end{array}$	D	C_5H_{12}
E	$\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$	F	C_3H_6

- 2.1 Define a *hydrocarbon*. (2)
- 2.2 Write down the letters that represent TWO unsaturated hydrocarbons. (2)
- 2.3 Write down the general formula of the following compounds:
- 2.3.1 **B** (1)
- 2.3.2 **F** (1)
- 2.4 Write down the IUPAC names of the following compounds:
- 2.4.1 **C** (2)
- 2.4.2 **D** (2)
- 2.5 Draw the structural formula of the following compounds:
- 2.5.1 **E** (2)
- 2.5.2 **F** (2)

[14] ...

QUESTION 3 (Start on a new page.)

The following isomers are commonly used as fuels because of their low boiling points:





- A** Butane
B 2-methylpropane

- 3.1 Identify the homologous series to which these compounds belong. (1)
- 3.2 Define the term *structural isomers*. (2)
- 3.3 What type of isomerism is represented by the compounds above? (1)
- 3.4 Identify the type of intermolecular forces in both compounds. (1)
- 3.5 Which ONE of the compounds has the strongest intermolecular forces?
Write down only **A** or **B**. (1)
- 3.6 Give a reason for the answer to QUESTION 3.5. (2)
- 3.7 Define *vapour pressure*. (2)
- 3.8 Write down the compound above with a higher:
- 3.8.1 Boiling point (1)
- 3.8.2 Vapour pressure (1)
- [12]**



QUESTION 4 (Start on a new page.)

Consider the following organic reactions.



4.1 Consider reaction 1.

4.1.1 Name the type of reaction. (1)

4.1.2 Write down the CONDENSED STRUCTURAL FORMULA of compound X. (2)

4.1.3 Is compound X a PRIMARY, SECONDARY or TERTIARY alcohol? (1)

4.2 Consider reaction 2.

4.2.1 Balance the chemical equation for this reaction. (1)

4.2.2 Is this an EXOTHERMIC or ENDOTHERMIC reaction? Give a reason. (3)

4.3 Consider reaction 3.

Product Y is used in cigarette lighters.

4.3.1 Name the type of ADDITION reaction taking place. (1)

4.3.2 Write down the IUPAC name of product Y. (2)

4.3.3 Write down the NAME or FORMULA of the catalyst used. (1)



4.4 Reaction 4 is a polymerisation reaction used to produce plastics.


4.4.1 Define the term *polymer*. (2)

4.4.2 Write down the NAME of the monomer used in this reaction. (1)
[15]



QUESTION 5 (Start on a new page.)

Scratches on chromium mag wheels are removed by electroplating. The diagram below represents an electrolytic cell involved in such a process.




- 5.1 Define the term *electrolytic cell*. (2)
- 5.2 Write down the NAME or FORMULA of the metal used as the anode. (1)
- 5.3 Which electrode, **A** or **B**, is the cathode? Give a reason. (2)
- 5.4 Write down the half reaction taking place at the following electrodes:
- 5.4.1 **A** (2)
- 5.4.2 **B** (2)
- 5.5 What is the purpose of the power source? (1)
- 5.6 Why is the DC source preferred to the AC source? (2)
- [12]**



QUESTION 6 (Start on a new page.)

The diagram below represents an electrochemical cell operating under standard conditions.



- 6.1 What type of reaction is taking place in the cell above? (1)
- 6.2 What is the reading on the voltmeter? (1)
- 6.3 Switch **S** is now closed.
 - 6.3.1 Calculate the reading on the voltmeter. (4)
 - 6.3.2 Which ONE of the electrodes, **Pb** or **Al**, will experience a decrease in mass? Explain the answer. (3)
 - 6.3.3 Write down the net ionic cell reaction. (3)

[12]

TOTAL: 75



**DATA FOR TECHNICAL SCIENCES GRADE 12
PAPER 2
GEGEWENS VIR TEGNIESE WETENSKAPPE GRAAD 12
VRAESTEL 2**

TABLE 1/TABEL 1: PHYSICAL CONSTANTS/FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,01 \times 10^5 \text{ Pa}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	$0 \text{ }^\circ\text{C}/273 \text{ K}$

TABLE 2/TABEL 2: FORMULAE/FORMULES

Emf/Emk	$E^\theta_{\text{cell}} = E^\theta_{\text{cathode}} - E^\theta_{\text{anode}} \quad / \quad E^\theta_{\text{sel}} = E^\theta_{\text{katode}} - E^\theta_{\text{anode}}$ or/of $E^\theta_{\text{cell}} = E^\theta_{\text{reduction}} - E^\theta_{\text{oxidation}} \quad / \quad E^\theta_{\text{sel}} = E^\theta_{\text{reduksie}} - E^\theta_{\text{oksidasie}}$ or/of $E^\theta_{\text{cell}} = E^\theta_{\text{oxidising agent}} - E^\theta_{\text{reducing agent}} \quad / \quad E^\theta_{\text{sel}} = E^\theta_{\text{oksideermiddel}} - E^\theta_{\text{reduseermiddel}}$
---------	---



TABLE 3: THE PERIODIC TABLE OF ELEMENTS / TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

KEY/SLEUTEL	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
	(I)	(II)		(IV)	(V)	(VI)	(VII)	(VIII)					(III)	(IV)	(V)	(VI)	(VII)	(VIII)	
Atomic number Atoomgetal	↓																		
1 1,01	1 H	2 4	3 Li	4 Be	5 9	6 12	7 Mg	8 24	9 20	10 Ca	11 40	12 38	13 88	14 56	15 137	16 88	17 88	18 226	19 89
11 0,9	11 Na	12 24	13 23	14 20	15 40	16 38	17 88	18 56	19 137	20 88	21 89	22 179	23 181	24 184	25 186	26 190	27 192	28 195	29 197
19 0,8	19 K	20 40	21 39	22 88	23 89	24 91	25 101	26 103	27 106	28 108	29 112	30 115	31 119	32 122	33 127	34 128	35 131	36 131	37 131
37 0,8	37 Rb	38 88	39 89	40 91	41 92	42 96	43 101	44 103	45 106	46 108	47 112	48 115	49 119	50 122	51 127	52 128	53 131	54 131	55 133
55 0,7	55 Cs	56 137	57 139	58 181	59 184	60 186	61 190	62 192	63 195	64 197	65 201	66 204	67 207	68 209	69 210	70 210	71 210	72 210	73 210
87 0,7	87 Fr	88 Ra	89 Ac	90 232	91 232	92 238	93 238	94 238	95 238	96 238	97 238	98 238	99 238	100 238	101 238	102 238	103 238	104 238	105 238
107 0,7	107 Uu	108 Uu	109 Uu	110 Uu	111 Uu	112 Uu	113 Uu	114 Uu	115 Uu	116 Uu	117 Uu	118 Uu	119 Uu	120 Uu	121 Uu	122 Uu	123 Uu	124 Uu	125 Uu
137 0,7	137 Nh	138 Nh	139 Nh	140 Nh	141 Nh	142 Nh	143 Nh	144 Nh	145 Nh	146 Nh	147 Nh	148 Nh	149 Nh	150 Nh	151 Nh	152 Nh	153 Nh	154 Nh	155 Nh
167 0,7	167 Lv	168 Lv	169 Lv	170 Lv	171 Lv	172 Lv	173 Lv	174 Lv	175 Lv	176 Lv	177 Lv	178 Lv	179 Lv	180 Lv	181 Lv	182 Lv	183 Lv	184 Lv	185 Lv
187 0,7	187 Og	188 Og	189 Og	190 Og	191 Og	192 Og	193 Og	194 Og	195 Og	196 Og	197 Og	198 Og	199 Og	200 Og	201 Og	202 Og	203 Og	204 Og	205 Og

Electronegativity Elektronegatiwiteit	↑																		
29 Cu	29 Cu	26 Fe	27 Co	24 Cr	25 Mn	22 Ti	23 V	20 Ca	21 Sc	19 K	18 Ar	17 Cl	16 S	15 P	14 Si	13 Al	12 C	11 B	10 Ne
63,5	63,5	56	59	52	55	48	51	40	45	39	36	35	32	31	28	27	16	14	10
Symbol Simbool	Cu	Fe	Co	Cr	Mn	Ti	V	Ca	Sc	K	Ar	Cl	S	P	Si	Al	C	B	Ne

Approximate relative atomic mass Benaderde relatiewe atoommassa	↑																		
29	29	56	59	52	55	48	51	40	45	39	36	35	32	31	28	27	16	14	10
63,5	63,5	56	59	52	55	48	51	40	45	39	36	35	32	31	28	27	16	14	10
Symbol	Cu	Fe	Co	Cr	Mn	Ti	V	Ca	Sc	K	Ar	Cl	S	P	Si	Al	C	B	Ne



TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE

Increasing strength of oxidising agents/Toenemende sterkte van oksideermiddels

Half-reactions/Halfreaksies		E° (V)
$F_2(g) + 2e^-$	$\rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^-$	$\rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^-$	$\rightleftharpoons 2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^-$	$\rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^-$	$\rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	$\rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^-$	$\rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^-$	$\rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^-$	$\rightleftharpoons Pt$	+ 1,20
$Br_2(l) + 2e^-$	$\rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^-$	$\rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^-$	$\rightleftharpoons Hg(l)$	+ 0,85
$Ag^+ + e^-$	$\rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^-$	$\rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^-$	$\rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^-$	$\rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^-$	$\rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^-$	$\rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^-$	$\rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^-$	$\rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^-$	$\rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^-$	$\rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^-$	$\rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^-$	$\rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^-$	$\rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^-$	$\rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^-$	$\rightleftharpoons Fe$	- 0,06
$Pb^{2+} + 2e^-$	$\rightleftharpoons Pb$	- 0,13
$Sn^{2+} + 2e^-$	$\rightleftharpoons Sn$	- 0,14
$Ni^{2+} + 2e^-$	$\rightleftharpoons Ni$	- 0,27
$Co^{2+} + 2e^-$	$\rightleftharpoons Co$	- 0,28
$Cd^{2+} + 2e^-$	$\rightleftharpoons Cd$	- 0,40
$Cr^{3+} + e^-$	$\rightleftharpoons Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^-$	$\rightleftharpoons Fe$	- 0,44
$Cr^{3+} + 3e^-$	$\rightleftharpoons Cr$	- 0,74
$Zn^{2+} + 2e^-$	$\rightleftharpoons Zn$	- 0,76
$2H_2O + 2e^-$	$\rightleftharpoons H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^-$	$\rightleftharpoons Cr$	- 0,91
$Mn^{2+} + 2e^-$	$\rightleftharpoons Mn$	- 1,18
$Al^{3+} + 3e^-$	$\rightleftharpoons Al$	- 1,66
$Mg^{2+} + 2e^-$	$\rightleftharpoons Mg$	- 2,36
$Na^+ + e^-$	$\rightleftharpoons Na$	- 2,71
$Ca^{2+} + 2e^-$	$\rightleftharpoons Ca$	- 2,87
$Sr^{2+} + 2e^-$	$\rightleftharpoons Sr$	- 2,89
$Ba^{2+} + 2e^-$	$\rightleftharpoons Ba$	- 2,90
$Cs^+ + e^-$	$\rightleftharpoons Cs$	- 2,92
$K^+ + e^-$	$\rightleftharpoons K$	- 2,93
$Li^+ + e^-$	$\rightleftharpoons Li$	- 3,05

Increasing strength of reducing agents/Toenemende sterkte van reduceermiddels



TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

Increasing strength of oxidising agents/Toenemende sterkte van oksideermiddels

Half-reactions/Halfreaksies		E ^θ (V)
Li ⁺ + e ⁻	⇌ Li	- 3,05
K ⁺ + e ⁻	⇌ K	- 2,93
Cs ⁺ + e ⁻	⇌ Cs	- 2,92
Ba ²⁺ + 2e ⁻	⇌ Ba	- 2,90
Sr ²⁺ + 2e ⁻	⇌ Sr	- 2,89
Ca ²⁺ + 2e ⁻	⇌ Ca	- 2,87
Na ⁺ + e ⁻	⇌ Na	- 2,71
Mg ²⁺ + 2e ⁻	⇌ Mg	- 2,36
Al ³⁺ + 3e ⁻	⇌ Al	- 1,66
Mn ²⁺ + 2e ⁻	⇌ Mn	- 1,18
Cr ²⁺ + 2e ⁻	⇌ Cr	- 0,91
2H ₂ O + 2e ⁻	⇌ H ₂ (g) + 2OH ⁻	- 0,83
Zn ²⁺ + 2e ⁻	⇌ Zn	- 0,76
Cr ³⁺ + 3e ⁻	⇌ Cr	- 0,74
Fe ²⁺ + 2e ⁻	⇌ Fe	- 0,44
Cr ³⁺ + e ⁻	⇌ Cr ²⁺	- 0,41
Cd ²⁺ + 2e ⁻	⇌ Cd	- 0,40
Co ²⁺ + 2e ⁻	⇌ Co	- 0,28
Ni ²⁺ + 2e ⁻	⇌ Ni	- 0,27
Sn ²⁺ + 2e ⁻	⇌ Sn	- 0,14
Pb ²⁺ + 2e ⁻	⇌ Pb	- 0,13
Fe ³⁺ + 3e ⁻	⇌ Fe	- 0,06
2H⁺ + 2e⁻	⇌ H₂(g)	0,00
S + 2H ⁺ + 2e ⁻	⇌ H ₂ S(g)	+ 0,14
Sn ⁴⁺ + 2e ⁻	⇌ Sn ²⁺	+ 0,15
Cu ²⁺ + e ⁻	⇌ Cu ⁺	+ 0,16
SO ₄ ²⁻ + 4H ⁺ + 2e ⁻	⇌ SO ₂ (g) + 2H ₂ O	+ 0,17
Cu ²⁺ + 2e ⁻	⇌ Cu	+ 0,34
2H ₂ O + O ₂ + 4e ⁻	⇌ 4OH ⁻	+ 0,40
SO ₂ + 4H ⁺ + 4e ⁻	⇌ S + 2H ₂ O	+ 0,45
Cu ⁺ + e ⁻	⇌ Cu	+ 0,52
I ₂ + 2e ⁻	⇌ 2I ⁻	+ 0,54
O ₂ (g) + 2H ⁺ + 2e ⁻	⇌ H ₂ O ₂	+ 0,68
Fe ³⁺ + e ⁻	⇌ Fe ²⁺	+ 0,77
NO ₃ ⁻ + 2H ⁺ + e ⁻	⇌ NO ₂ (g) + H ₂ O	+ 0,80
Ag ⁺ + e ⁻	⇌ Ag	+ 0,80
Hg ²⁺ + 2e ⁻	⇌ Hg(l)	+ 0,85
NO ₃ ⁻ + 4H ⁺ + 3e ⁻	⇌ NO(g) + 2H ₂ O	+ 0,96
Br ₂ (l) + 2e ⁻	⇌ 2Br ⁻	+ 1,07
Pt ²⁺ + 2 e ⁻	⇌ Pt	+ 1,20
MnO ₂ + 4H ⁺ + 2e ⁻	⇌ Mn ²⁺ + 2H ₂ O	+ 1,23
O ₂ (g) + 4H ⁺ + 4e ⁻	⇌ 2H ₂ O	+ 1,23
Cr ₂ O ₇ ²⁻ + 14H ⁺ + 6e ⁻	⇌ 2Cr ³⁺ + 7H ₂ O	+ 1,33
Cl ₂ (g) + 2e ⁻	⇌ 2Cl ⁻	+ 1,36
MnO ₄ ⁻ + 8H ⁺ + 5e ⁻	⇌ Mn ²⁺ + 4H ₂ O	+ 1,51
H ₂ O ₂ + 2H ⁺ + 2 e ⁻	⇌ 2H ₂ O	+1,77
Co ³⁺ + e ⁻	⇌ Co ²⁺	+ 1,81
F ₂ (g) + 2e ⁻	⇌ 2F ⁻	+ 2,87

Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels



